

## The role of environmental feedback in a brain state switch from passive to active sensing

Article (Published Version)

Buckley, Christopher L and Toyoizumi, Taro (2013) The role of environmental feedback in a brain state switch from passive to active sensing. BMC Neuroscience, 14 (Supp 1). P395. ISSN 1471-2202

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/59443/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

POSTER PRESENTATION

Open Access

# The role of environmental feedback in a brain state switch from passive to active sensing

Christopher L Buckley\*, Taro Toyozumi

From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013  
Paris, France. 13-18 July 2013

Coherent behaviour emerges from mutual interaction between the brain, body and environment across multiple timescales and not from within the brain alone [1,2]. For example sensation is actively shaped by dynamical interaction of the brain and environment through motor actions such as sniffing, saccading, and touching. The onset of active sensing is often concomitant with qualitative changes in neural dynamics [3] and responses to sensory input [4,5]. Indeed, some neural responses are uniquely sensitive to the presence or absence of dynamical sensory feedback [6]. However, understanding how active sensing strategies impact on neural dynamics and sensory responses is an open challenge. Whisking behaviour in rodents has been the central model system for studying neural mechanisms of active sensing. During quiet wakefulness the membrane potential of neurons in the barrel cortex exhibit high power, low frequency fluctuations and nearby neurons become highly correlated [3]. As active whisking onsets, the brain state of the barrel cortex qualitatively changes: low frequency fluctuations are suppressed and nearby neurons decorrelate [3]. Interestingly both correlation and low frequency LFP are partially restored during periods of active touch, i.e., when the mouse palpate towards, and repeatedly contacts with an object [3]. The brain state transition is also concomitant with changes in responses of barrel cortex neurons to whisker stimulation. The sensitivity of neurons to whisker perturbation drops with the onset of whisking [5]. However, robust and repeatable whisker response are recovered for more naturalistic stimuli, i.e., active touch [5].

Here we propose a theory, and construct a model, of active whisking that explains the core phenomenology through the dynamical interaction between the brain

and the environment. The three core assumptions underlying the theory are:

1. Strong low frequency membrane fluctuations, intra-neural correlations between nearby neurons, and sensory responses to brief whisker deflection associative with quiet attentive state arise as result of network dynamics that are close to a dynamical instability.
2. Reafferent input (sensory feedback related to self-action) during whisking behaviour provides negative feedback to sensory neurons that stabilize cortical dynamics reducing low frequency fluctuations, intra neural correlations, sensory responses to brief whisker deflection, while increasing the correlation between cortical activity and whisker position.
3. Interrupting the reafferent signal, via a whisker touch event, temporally destabilise the cortex and enhances ex-afferent input (external sensory input coding whisker contacts).

Our theory and model suggests that sensory feedback via reafference signals is *sufficient* to account for the changes in cortical dynamics that appear with the onset of active whisking and provides a novel mechanistic account for sensory processing during active whisking. We discuss how this feedback stabilization mechanism coexists and interacts with other internal mechanisms that modulate cortical dynamics within the brain [3,7]. We quantify how this mechanism of active touch enhances the signal-to noise ratio of the input to the cortex. Finally we discuss the wider implications of these results for experimental work attempting to characterise neural activity and responses in an open-loop (in the absence of sensory/motor feedback) condition.

\* Correspondence: [chrisbuckley@brain.riken.jp](mailto:chrisbuckley@brain.riken.jp)  
Lab for Neural Computation and Adaptation, Riken Brain Science Institute,  
Tokyo, Japan

Published: 8 July 2013

## References

1. Gibson JJ: **Observations on active touch.** *Psychological Review* 1962, **69**(6):477-491.
2. Beer RD: **Dynamical approaches to cognitive science.** *Trends in Cognitive Science* 2000, **4**(3):91-99.
3. Poulet JFA, Petersen CCH: **Internal brain state regulates membrane potential synchrony in barrel cortex of behaving mice.** *Nature* 2008, **454**(7206):881.
4. Otazu GH, Tai LH, Yang Y, Zador AM: **Engaging in an auditory task suppresses responses in auditory cortex.** *Nature Neuroscience* 2009, **12**(5):646-654.
5. Crochet S, Petersen CCH: **Correlating whisker behavior with membrane potential in barrel cortex of awake mice.** *Nature Neuroscience* 2006, **9**(5):608-610.
6. Keller GB, Bonhoeer T, and Hubener M: **Sensory/motor mismatch signals in primary visual cortex of the behaving mouse.** *Neuron* 2012, **74**(5):809-815.
7. Poulet JFA, Fernandez LMJ, Crochet S, Petersen CCH: **Thalamic control of cortical states.** *Nature Neuroscience* 2012, **15**(3):370-372.

doi:10.1186/1471-2202-14-S1-P395

**Cite this article as:** Buckley and Toyoizumi: The role of environmental feedback in a brain state switch from passive to active sensing. *BMC Neuroscience* 2013 **14**(Suppl 1):P395.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

